

Gravitational field of an infinitely long supermassive cosmic string

M. M. Som

Instituto de Física

Universidade Federal do Rio de Janeiro

Rio de Janeiro, Brasil

and

Marcelo de Oliveira Souza

Universidade Estadual do Norte Fluminense

Centro de Ciência e Tecnologia

Campos, Rio de Janeiro, Brasil

and

Instituto de Física

Universidade Federal do Rio de Janeiro

Rio de Janeiro, Brasil

PACS number(s): 04.20.Jb, 04.20.Cv

e-mail: som@if.ufrj.br and mm@uenf.br

11/11/94

Abstract

We obtain an exact solution of the coupled Einstein-scalar-gauge field equations for a local infinitely long supermassive cosmic string. The solution corresponds to that of Hiscock-Gott. The string appears to be due to the freezing of the scalar field at the null value giving rise to a constant linear energy density.

Spontaneous symmetry breaking in gauge theories leads to phase transitions in the early Universe [1,2,3]. In this period can appear topological structures in the Universe like strings, vacuum domain walls and monopoles. One of these topological defects that has drawn more attention is the cosmic string. The cosmic strings are linear topological defects. Only infinitely long and closed loop strings can exist. An infinitely long cosmic string is a static cylindrically symmetric configuration of self-interacting scalar field minimally coupled to a U(1) gauge field.

We have two general type of strings. Those which arise in a phase transition in which a gauge symmetry undergoes spontaneous symmetry breaking is called local or gauge strings, and those which we call a global string, arise as a result of a spontaneous breaking of global symmetry. Strings that have cosmological interest are the local strings.

Local strings have a linear energy density $\mu \sim \eta^2$, where η is the vacuum expectation value (VEV) acquired by the gauge field in the phase transition.

Strings of astrophysical relevance were formed during phase transitions at the grand-unified-theory (GUT) scale (10^{15} GeV). They have linear energy density $\mu \sim \eta^2$ of the order 10^{-6} . This small value of μ could justify the weak-field approximation used by Vilenkin.

Vilenkin [4] using the weak-field approximation obtained that the geometry around a cosmic string is conical, with the deficit angle given by $\delta\phi = 8\pi\mu$, where μ is the linear energy density of the string. Hiscock [5] and Gott [6] found independently an exact spacetime metric corresponding to a static cylindrically symmetric string, and showed that the results obtained by Vilenkin are correct. They utilized a stress energy tensor with the energy density constant within the string, without considering the structure of the scalar and the gauge fields. Raychaudhuri [7] criticized these results giving a new interpretation to their metric, using the full equations, not taken into account by them. So far no solution in closed form is obtained of the coupled Einstein-scalar-gauge field equations.

More massive strings, $\eta \gg 10^{-3}$ could be produced, because the Universe might have undergone phase transitions at energy scales higher than the 10^{15} GeV GUT scale. Due to their large masses, $\mu \gg 10^{-6}$, we believe it is not reasonable to treat these strings by means of the weak-field approximation.

We are going to consider in this paper supermassive strings. They are strings with $\eta \gg 10^{-3}$.

The model of the self-interacting potential for the scalar complex field is provided by the standard "Mexican hat" potential. Then, the only consistent way to find the metric describing an infinitely long cosmic string is to solve the coupled Einstein-scalar-gauge equations.

We consider the simplest case of a local U(1) symmetry. After spontaneous symmetry breaking, the manifold of degenerate vacua is a circle; since this manifold is not simply

connected, strings can occur. The simplest scalar gauge field theory leading to the formation of cosmic strings is the Abelian Higgs model which contains a U(1) gauge field A_μ and a complex charged scalar field $\Phi = R \exp(i\psi)$ with a coupling constant e and a potential

$$V(\Phi) = \frac{\lambda}{2}(\Phi\Phi^* - \eta^2) \quad (1)$$

where λ and η are two coupling constants. (We use units where $G = c = 1$.)

The Lagrangian for these fields is,

$$L = -\frac{1}{2}\nabla^\mu R \nabla_\mu R - \frac{1}{2}R^2(\nabla_\mu \psi + eA_\mu)(\nabla^\mu \psi + eA^\mu) - \lambda(R^2 - \eta^2)^2 - \frac{1}{16\pi}F_{\mu\nu}F^{\mu\nu}. \quad (2)$$

where $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu$.

The complex scalar field and the components of the gauge field take the form

$$\begin{aligned} R &= R(r) \quad , \quad \psi = n\phi, \\ A_\mu &= \frac{1}{e}[P(r) - n]\nabla_\mu \phi, \end{aligned} \quad (3)$$

where n is the winding number.

The spacetime metric assumed to be static and cylindrically symmetric, as shown by Shaver [8], is given by,

$$ds^2 = -dt^2 + dr^2 + dz^2 + \beta(r)^2 d\phi^2 \quad (4)$$

with the angular coordinate ϕ ranges from 0 to 2π as usual.

To get solutions with the metric regularity at $r = 0$, and with a finite energy, we take the boundary conditions

$$\begin{aligned} \lim_{r \rightarrow 0} R(r) &= 0 \quad \text{and} \quad \lim_{r \rightarrow 0} P(r) = n \\ \lim_{r \rightarrow \infty} R(r) &= \eta \quad \text{and} \quad \lim_{r \rightarrow \infty} P(r) = 0. \end{aligned} \quad (5)$$

Linet [9] proved the Bogomol'nyi inequality on the linear mass density μ of these solutions in the case $e^2 \leq 8\lambda$. The lower bound $\mu = \pi|n|\eta^2$ correspond a solution with $e^2 = 8\lambda$ and the metric has necessarily the form (4). We are going to consider the case $n = 1$. On the axis $r = 0$ the function β has to satisfy the following condition

$$\lim_{r \rightarrow 0} \frac{\beta^2}{r^2} = 1. \quad (6)$$

In this case the non-vanishing components of G_ν^μ are

$$G_z^z = G_t^t = \frac{1}{\beta} \frac{d^2\beta}{dr^2}. \quad (7)$$

The components of the energy-momentum tensor are

$$T_t^t = T_z^z = -\frac{1}{2}[(R')^2 + \frac{R^2 P^2}{\beta^2} + 2\lambda(R^2 - \eta^2)^2 + \frac{(P')^2}{\beta^2 e^2}] \quad (8)$$

$$T_r^r = \frac{1}{2}[(R')^2 - \frac{R^2 P^2}{\beta^2} - 2\lambda(R^2 - \eta^2)^2 + \frac{(P')^2}{\beta^2 e^2}] \quad (9)$$

$$T_\phi^\phi = \frac{1}{2}[-(R')^2 + \frac{R^2 P^2}{\beta^2} + 2\lambda(R^2 - \eta^2)^2 + \frac{(P')^2}{\beta^2 e^2}]. \quad (10)$$

When we consider $e^2 = 8\lambda$, the function R and P must obey the following Bogomol'nyi equations

$$P' = \frac{\beta e^2}{2}(R^2 - \eta^2) \quad (11)$$

$$R' = \frac{RP}{\beta} \quad (12)$$

and the Einstein equations reduce to,

$$\frac{\beta''}{\beta} = -8\pi(\frac{R^2 P^2}{\beta^2} + \frac{e^2}{4}(R^2 - \eta^2)) \quad (13)$$

Using (11) and (12) in (13), we can evaluate the integration, obtaining the following equation

$$\beta' = -4\pi(R^2 - \eta^2)P + 1 - 4\pi\eta^2. \quad (14)$$

Making the transformation

$$\beta = \frac{2\sqrt{2}\bar{\beta}}{\eta e}, \quad (15)$$

$$R = \eta \bar{R}, \quad (16)$$

and

$$r = \frac{2\sqrt{2}\bar{r}}{\eta e}, \quad (17)$$

as proposed by Shaver[8] , we see that the equations (11) and (14) become,

$$\bar{P}' = 4\bar{\beta}(\bar{R}^2 - 1) , \quad (18)$$

$$\bar{\beta}' = -4\pi\eta^2(\bar{R}^2 - 1)\bar{P} + 1 - 4\pi\eta^2 \quad (19)$$

and the equation (12) remains the same.

The set of equations (12), (18-19) with the boundary conditions (5) describe the cosmic strings. However, the exact solutions of the set of equations are so far not obtained. Linet [10] attempted to obtain exact solutions for the special case of a supermassive local string $4\pi\eta^2 = 1$.

For a supermassive local string, we consider $4\pi\eta^2 = 1$, then the equation (19) reduces to,

$$\bar{\beta}' = -(\bar{R}^2 - 1)\bar{P}. \quad (20)$$

From the equations (20) and (18) we obtain that,

$$\bar{P} = \sqrt{1 - 4\bar{\beta}^2} , \quad (21)$$

and,

$$2\frac{\bar{R}'}{\bar{R}} = \frac{(\frac{-\bar{\beta}'}{\bar{P}})'}{1 - \frac{\bar{\beta}'}{\bar{P}}} \quad (22)$$

Substituting (21) and (22) in (12) , we construct the following differential equations for $\bar{\beta}$

$$4\bar{\beta}^2(\bar{\beta}')^2 + (1 - 4\bar{\beta}^2)\bar{\beta}\bar{\beta}'' + 2(1 - 4\bar{\beta}^2)^2 - 2(1 - 4\bar{\beta}^2)^{\frac{3}{2}}\bar{\beta}' = 0. \quad (23)$$

An exact solution of the differential equation is given by,

$$\bar{\beta} = \frac{1}{2} \sin(2\bar{r}). \quad (24)$$

Then we have ,

$$\bar{P} = \cos(2\bar{r}). \quad (25)$$

Substituting (16) and (17) in (24) and (25), we have

$$\beta = \frac{1}{a} \sin(ar) , \quad (26)$$

$$P = \cos(ar) , \quad (27)$$

where $a = \frac{e}{8\sqrt{2\pi}}$.

We conclude that, in this case with a constant energy momentum tensor as the source, $R = 0$, what leads us to consider $\phi = 0$. Hiscock[5] obtained the same solution.

Recently Novello and Da Silva [11] obtained the exact solution in this form for the gravitational field of string related to the fundamental states of bosons of arbitrary mass.

To obtain the solution in the exterior region of the string we put $T_\mu^\nu = 0$. From (8-10) one obtains $R = \eta$ and $P = 0$.

The solution of the equation (13) is then given by $\beta = br + c$, where b and c are constants. The constant c can be removed by shifting the origin, the metric then takes the form

$$ds^2 = -dt^2 + dr^2 + dz^2 + b^2 r^2 d\phi^2. \quad (28)$$

In this metric, that is Minkowski flat, the angular coordinate ϕ ranges from 0 to $2\pi b$. Applying the usual Darmois-Lichnerowicz junction conditions (in this case reduces to $g_{\phi\phi}^+ = g_{\phi\phi}^-$ and $\frac{dg_{\phi\phi}^+}{dr} = \frac{dg_{\phi\phi}^-}{dr}$, where $+$ represent the exterior metric and $-$ the interior) at the boundary $r = r_0$, we find that

$$\frac{1}{a} \sin(ar_0) = br_0, \quad (29)$$

and,

$$\cos(ar_0) = b. \quad (30)$$

The same results can be obtained from the junction conditions used by Hiscock [5]: $g_{\mu\nu}^+ = g_{\mu\nu}^-$ and $K_{\mu\nu}^+ = K_{\mu\nu}^-$, where $K_{\mu\nu}$ is the extrinsic curvature.

Evaluating the linear energy density we obtain,

$$\begin{aligned} \mu &= \int_0^{r_0} \int_0^{2\pi} T_t^t \beta d\phi dr \\ &= \frac{1}{4} [1 - \cos(ar_0)]. \end{aligned} \quad (31)$$

From the equations (30) and (31), we get

$$b = (1 - 4\mu) \quad (32)$$

The exterior metric takes the form

$$ds^2 = -dt^2 + dz^2 + dr^2 + (1 - 4\mu)^2 r^2 d\phi^2. \quad (33)$$

Here we can define an angular coordinate $\phi' = (1 - 4\mu)\phi$ that ranges from 0 to $2\pi(1 - 4\mu)$, giving a conical angular deficit $\delta\phi = 8\pi\mu$.

From (8-10) one finds that,

$$T_t^t(\phi = 0) = T_t^t(A_\phi) = 2\lambda\eta^4 \quad (34)$$

in the interior region of the string.

Since the ϕ -field acquires the null value in the string core as shown by Linde[12] during the inflationary period we may consider that after the phase transition the ϕ -field settles down to the null value producing constant energy density. The gauge field also contributes the same amount of constant energy density.

However it should be remarked that this configuration of the string is possible only in the case of a supermassive local strings with $4\pi\eta^2 = 1$.

The authors would like to thank CNPq and UENF for a grant.

References

- [1] D.A. Kirzhnits, JETP Lett. 15, 745 (1972).
- [2] A.D. Linde, Rep. Prog. Phys. 42, 389 (1979).
- [3] T.W. Kibble, J. Phys. A9, 1387 (1976).
- [4] A. Vilenkin, Phys. Rev. D23, 852 (1981).
- [5] W. Hiscock, Phys. Rev. D31, 3288 (1985).
- [6] J. R. Gott III, Astrophys. J. 288, 422 (1985).
- [7] A. K. Raychauduri, Phys. Rev. D41, 3041 (1990).
- [8] E. Shaver, GRG 24, 187 (1992).
- [9] E. B. Linet, Phys. Lett. 124A, 240 (1990).
- [10] E. B. Linet, Class. Quantum Grav. 7, L75 (1990).
- [11] M. Novello and M.C. Motta da Silva, Phys. Rev. D48, 5017 (1993).
- [12] A.D. Linde, Phys. Rev. D50, 2456 (1994).